CLINICS IN PODIATRIC MEDICINE AND SURGERY

Clin Podiatr Med Surg
20 (2003) 803–824

Cumulative Index 2003

Volume 20

January	EXTERNAL FIXATION OF THE FOOT AND ANKLE, pages 1–198
April	PERIOPERATIVE MANAGEMENT OF THE PODIATRIC PATIENT, pages 199–372
July	CLINICAL PODOGERIATRICS: ASSESSMENT, EDUCATION AND PREVENTION, pages 373–634
October	DIABETES MELLITUS, pages 635–824

Note: Page numbers of article titles are in **boldface** type.

A

Abductor digiti minimi flap, in diabetic foot reconstruction, 779

Abductor hallucis flap, in diabetic foot reconstruction, 778

Ablative surgery, in diabetic foot, 720–721

ABLE (Althof-Beasley Lower Extremity Presurgical Assessment Tool), for psychological management, 204–208

Abscess, subungual, 432

Acarbose, for diabetes mellitus, 221–222, 643

Accidents, risk assessment for, in long-term care, 392

Acetaminophen, for diabetic neuropathy, 679

Acetohexamide, for diabetes mellitus, 640–641

Achilles tendon
- anatomy of, 362
- insertion tendinosis of, retrocalcaneal exostectomy for, **361–366**
- lengthening of, in diabetes mellitus, 790

Advanced glycation end products, in diabetes mellitus, 677

Age considerations, in external fixation, 22

Akin osteotomy, external fixation for, 35–36

Allopurinol, for gout, 483

Alpha-glucosidase inhibitors, for diabetes mellitus, 221–222, 643

Alpha-methyl dopa, preoperative evaluation of, 225

Alternative medications, preoperative evaluation of, 232–234

Althof-Beasley Lower Extremity Presurgical Assessment Tool, for psychological management, 204–208

Amantadine, for diabetic neuropathy, 683

American Society of Anesthesiologists, surgical risk categories of, 616–617

Amikacin, for diabetic foot infections, 664

Amitriptyline, for diabetic neuropathy, 680–681

Amoxicillin/clavulanate, for diabetic foot infections, 659–660

Ampicillin, for diabetic foot infections, 664

Ampicillin/sulbactam, for diabetic foot infections, 659–660, 662–664

Amputation
- in Charcot foot, 747
- in diabetic foot, 399, 611–612, 720–721, **793–801**
  - comorbidity assessment in, 795–796
  - debridement with, 796
  - dysvascular tissue removal in, 796–797
  - energy expenditures after, 799
  - epidemiology of, 710
  - in comprehensive treatment plan, 794–795

0891-8422/03/$ – see front matter © 2003 Elsevier Inc. All rights reserved.
doi:10.1016/S0891-8422(03)00101-0

in infections, 665
in limb salvage, 794–797
incision planning in, 797
indications for, 793
prevention of. *See* Surgery, for diabetic foot, elective.
racial factors in, 799–800
rethinking, 797
rigid deformities and, 795
risk factors for, 783–784
statistics of, 793–794
Syme, 797–798
in severe trauma, versus salvage, 301

Analgesia, perioperative
epidural, cryotherapy effects on, 263–264
patient-controlled, 262
preemptive, **237–255**
spinal, 263–264

Anesthesia
for extracorporeal shock wave therapy, for plantar fasciitis, 329
for trauma repair, 300–301

Angiography, in arterial occlusive disease, in diabetes mellitus, 699–702

Angioplasty, of peripheral vessels, in diabetes mellitus, 702–704

Angiotensin-converting enzyme inhibitors
for diabetic neuropathy, 683
preoperative evaluation of, 226

Ankle
arthrodesis of, in degenerative disease, 345–346
Charcot. *See* Charcot foot and ankle.
external fixation applications in, **65–96**
arthrodesis, 164–169
tibiocalcaneal, 16, 85–89
tibiotalar, 16, 74–85
triple, 16, 89–90
biomechanics of, 67–69
capabilities of, 66
circular, 67–69
complications of, 92–93
distraction arthrodiastasis, 90–92
frame removal in, 94
limitations of, 66–67
wire considerations in, 69–74

Ankle block, in trauma, 298

Ankle joint replacement, case review of, **335–359**
discussion, 342–349, 357
history of injury, 336–337, 350–352
physical examination, 337–339
postoperative course, 340–342, 355–356
procedure, 339–340

Anonychia, 431

Antiarrhythmic agents, for diabetic neuropathy, 682

Antibiotics, for diabetic foot infections, 658–665, 719

Anticoagulants
for deep venous thrombosis prophylaxis, 277–279
preoperative evaluation of, 226–229

Anticonvulsants, for diabetic neuropathy, 681–682

Antidepressants
for diabetic neuropathy, 680–681
preoperative evaluation of, 231

Antiplatelet agents, preoperative evaluation of, 230

Antipsychotics, preoperative evaluation of, 231

Anxiety, foot pain in, 401

Anxiolytics, preoperative evaluation of, 231

AO/ASIF system, for subtalar arthrodesis. *See* Synthes tubular external fixation.

Argatroban, for deep venous thrombosis, 278–279

Arterial occlusive disease
in diabetes mellitus, **689–708**
amputation risk in, 784
arterial reconstruction in, 702–704
evaluation of
angiography in, 699–701
Doppler studies in, 697–699
exercise testing in, 696
general assessment in, 690–691
limb pressure measurements in, 693–695
magnetic resonance imaging in, 701–702
physical examination in, 691–693
pulse volume recordings in, 695–696
toe pressure measurements in, 695
transcutaneous oximetry in, 696–697
pathophysiology of, 689
treatment of, 702–704, 719
in elderly persons
healing ability and, 531–532
history of, 527–532
noninvasive evaluation of, 539–543

physical examination in, 532–538
risk factors for, 530–531

Arterial ulcers, in elderly persons, 389, 399

Arthritis
degenerative. *See* Osteoarthritis.
in elderly persons, **469–485**
biomechanics of, 470–472
diagnosis of, 378
epidemiology of, 469–470
flatfoot in, 502
gout, 481–483
hallux rigidus in, 491–493
osteoarthritis. *See under* Osteoarthritis.
psoriatic, 462–463
reactive versus nonreactive joints in, 472–473
rheumatoid, 477–481
treatment of, 472–474
types of, 469
rheumatoid. *See* Rheumatoid arthritis.

Arthrodesis
ankle, in degenerative disease, 345–346
external fixation for
ankle, 164–169
calcaneocuboid, 54–56
first metatarsocuneiform, 33–35
first metatarsophalangeal, 30–32
naviculocuneiform, 57–58
subtalar, **181–194**
talonavicular, 56–57
tarsometatarsal, 52–54
tibiocalcaneal, 16, 85–89
tibiotalar, 16, 74–85
triple, 16, 89–90
for Charcot foot, 751–752

Arthrodiastasis, distraction, 90–92

Arthrometric examination, of deformities, 514–515

Arthropathy, Charcot. *See* Charcot foot and ankle.

Arthroplasty, hallux, in diabetes mellitus, 786–787

Arthrosis, post-traumatic, ankle joint replacement for, case review of, **335–359**

Aspirin
for deep venous thrombosis, 282
preoperative evaluation of, 230–232

Assessment, of elderly persons, 378–379, **407–429**
clinical findings, 413–414
for accident risk, in long-term care, 392
for biomechanical conditions, 417–418
for chronic diseases, 411–412
for diabetes mellitus, **573–582**
for pressure ulcers, 384–385
in long-term care, 383–389
initial, 408–414
Medicare class findings in, 414–415
Medicare codes and, 408–411
neurological, **547–571**
outline for, 416–417
preoperative, 619–621
protocol for, 418–424
symptoms, 412–413
systems review in, 408

Association for the Study of Internal Fixation (AO/ASIF), fracture management principles of, 292–293

Atenolol, preoperative evaluation of, 225

Atherosclerosis, in peripheral vascular disease, 530, 690–691

Autonomic dysfunction, assessment of, in diabetes mellitus, 713

Avulsion, of nail, 434

Aztreonam, for diabetic foot infections, 664

B

Babinski's sign, in neurological evaluation, 567

Balance realignment therapy, for deformities, 521

Beau's lines, 436

Becaplermin, for diabetic ulcers, 717

Benzodiazepines
for diabetic neuropathy, 679
preoperative evaluation of, 231

Beta-blockers, preoperative evaluation of, 224–225

Beta-lactamase inhibitors, for diabetic foot infections, 663

Betacarotene, for warts, 464

Biguanides, for diabetes mellitus, 222, 641–642

Bilobed flaps, in diabetic foot reconstruction, 767–769

Biomechanics
of arthritis, 470–472
of deformities, evaluation of, 512–518
of diabetic foot, 575–576
of hyperkeratosis formation, 444–445

Biopsy
of bone, in osteomyelitis, in diabetes mellitus, 658–659
of skin lesions, 454–456

Bisoprolol, preoperative evaluation of, 225

Blisters, 385, 464–465

Block balance testing, for deformities, 521–523

Blood vessels, injury of, in external fixation, 92

Bone
biopsy of, in osteomyelitis, in diabetes mellitus, 658–659
destructive changes of, in diabetes mellitus. *See* Charcot foot and ankle.

Bone block distraction, in calcaneocuboid joint arthrodesis, 54–56

Bone graft, external fixation for, 36

Bone scan, in diabetic foot, 732–735
with osteoarthropathy, 657
with ulcers, 714–715

Brachymetatarsal, correction of, external fixation for, 36–38

Braden Scale, for pressure ulcer staging, 385

Brief Pain Inventory, 258

Buechel-Pappas ankle prosthesis, 339–340, 348–349, 351–356

Bullous diseases, 464–465

Bunion. *See* Hallux valgus.

Bunionette (tailor's bunion), 497–498

Bupivacaine, for preemptive analgesia, 246

Bursitis, retrocalcaneal, retrocalcaneal exostectomy for, **361–366**

C

Calcaneocuboid joint, arthrodesis of, external fixation for, 54–56

Calcaneus
fractures of, external fixation for, 169–170
spurs on, retrocalcaneal exostectomy for, **361–366**

Callus (tyloma), 397–398, 445

Callus distraction, external fixation for, 19

Cancer, skin, in elderly persons, 463–464

Capsaicin
for diabetic neuropathy, 682–683
for osteoarthritis, 476

Carbamazepine, for diabetic neuropathy, 681

Cardiovascular drugs, preoperative evaluation of, 224–226

Cardiovascular system, preoperative evaluation of, 214

Cast
for Charcot foot, 747
total-contact, for diabetic ulcers, 718
versus external fixation, 13

Cavovarus foot, external fixation for, 49

Cavus foot, 503–505
midfoot osteotomy for, external fixation for, 58–59

Cefadroxil, for diabetic foot infections, 659–660

Cefazolin, for diabetic foot infections, 658

Cefepime, for diabetic foot infections, 659, 662, 664

Cefoperazone, for diabetic foot infections, 664

Cefotaxime, for diabetic foot infections, 664

Cefoxitin, for diabetic foot infections, 662–663

Ceftazidime, for diabetic foot infections, 662–664

Ceftriaxone, for diabetic foot infections, 662–663

Celecoxib, for perioperative pain, preemptive administration of, 250

Cephalexin, for diabetic foot infections, 659

Cerebellar function, assessment of, 568–569

Charcot foot and ankle, **97–117, 741–756**
acute
diagnosis of, 746
treatment of, 747–751
chronic, treatment of, 751–754
classification of, 730, 744–746
clinical features of, 730
definition of, 97
diagnosis of, 746
differential diagnosis of, 730
etiology of, 741–742
external fixation for, 19, 49–52, 59–60, 747–753
goals of, 98–100
postoperative care in, 112–116
surgical options in, 101–103
techniques for, 104–112

imaging in, 656–658, **729–740**
in diabetes mellitus, 578
incidence of, 97–98
pathophysiology of, 729–730, 742–743
stages of, 98
treatment of, 743–744, 747, 789–790

Charnley, J.C., external fixation devices of, 5

Charnley classification, of foot and ankle trauma, 298

Chlorpropamide, for diabetes mellitus, 640–641

Ciprofloxacin, for diabetic foot infections, 659–660, 662, 664–665

Claudication, intermittent, in peripheral vascular disease, 527–529, 690

Claw toes, 489–491

Clinafloxacin, for diabetic foot infections, 660

Clindamycin, for diabetic foot infections, 659–660, 662, 664–665

Clonidine
for diabetic neuropathy, 683
preoperative evaluation of, 225

Clopidogrel, preoperative evaluation of, 230

Closed reduction, in Charcot foot, 749, 751

Clots, deep venous. *See* Deep venous thrombosis.

Clubbing, of nails, 432

Clubfoot, correction of, external fixation for, 17–18

Coagulation disorders
deep venous thrombosis in, 271
from alternative medications, 232–233

Colchicine, for gout, 482

Cold therapy. *See* Cryotherapy.

Collagen dressing, for diabetic ulcers, 717

Coma, in diabetes mellitus, 639

Compartment syndrome, in ankle trauma, 295–296

Complementary medications, preoperative evaluation of, 232–234

Comprehensive care plans, for nursing home residents, 389

Compression, in forefoot applications, external fixation for, 30–36

Computed tomography
in deep venous thrombosis, 275, 280
in diabetic foot, 737
in diabetic foots, with osteoarthropathy, 657–658

Computer-assisted sensory examination, in diabetes mellitus, 675

Computerized blood flow studies, in diabetic foot, 734

Confusion, postoperative, 623

Congenital nail disorders, 432

Contact dermatitis, 459–460

Contractures, in trauma, external fixation for, 174, 176

Contusions, of nail plate, 434

Corn
hard (heloma durum), 445
nail bed, 434, 445
seed (heloma miliare), 445
soft (heloma molle), 446

Coronary artery disease, peripheral vascular disease and, 530

Corticosteroids
for skin disorders, 461–462
intraarticular
for gout, 482
for osteoarthritis, 477
for rheumatoid arthritis, 480
preoperative evaluation of, 230

Cranial neuropathy, in diabetes mellitus, 674

Cryotherapy, postoperative, **307–322**
continuous-flow
case reports on, 311–314
normal subject trials of, 314–319
prescription for, 319–320
injury from, 310–311
pain physiology and, 307–310

Culture, in infections, in diabetes mellitus, 656, 719

Curative surgery, in diabetic foot, 720

Custom-molded footwear, for elderly persons, 593, 595, 598, 600

Cyclooxygenase inhibitors, for perioperative pain, 250, 260

D

D-dimer degradation products, in deep venous thrombosis, 275–276

Dalteparin, for deep venous thrombosis, 278

Debridement
in trauma, 299–300, 302

of diabetic foot infections, 665, 716
of pressure ulcers, 386

Deep venous thrombosis, in surgery, **269–289**
anatomic considerations in, 272
case reviews of, 283–285
diagnosis of, 274–276
epidemiology of, 269
for trauma, 300, 304–305
natural history of, 271–272
pathophysiology of, 270–272
prophylaxis for, 228–229, 276–277, 281–283, 300
risk factors for, 269–270, 272–273, 281
treatment of, 277–280

Deformities
external fixation correction of, 16–18
in midfoot, 47–49
lower leg. *See* Lower leg deformities.
in diabetes mellitus
elective reconstruction of, 720–721
surgical treatment of. *See* Surgery, for diabetic foot, elective.
treatment of, 795
ulcers in, 711–713, 720–721
in elderly persons, **487–509, 511–526**
biomechanical evaluation of, 512–518
claw toes, 489–491
compensatory malalignments in, 518–519
footwear for, 523–526, 595–596. *See also* Footwear.
hallux rigidus, 491–493
hallux valgus. *See* Hallux valgus.
hammertoes, 487–491
heel pain, 505–508
in diabetes mellitus, 575
in osteoarthritis, 475–476
metatarsalgia, 498–501
orthoses for, 523–526. *See also* Orthoses.
pain in, 398
pathomechanics of, 518–523
pes cavus, 503–505
pes planus, 501–503
surgery for, 610–611
tailor's bunion, 497–498

Delirium, postoperative, 623

Depression, surgical outcome and, 202

Dermatitis, contact, 459–460

Dermatitis herpetiformis, 465

Dermis, tissue-engineered, for diabetic ulcers, 717–718

Desipramine, for diabetic neuropathy, 680

Diabetes mellitus, **635–653**
amputation in. *See* Amputation, in diabetic foot.
Charcot foot and ankle in. *See* Charcot foot and ankle.
classification of, 636
diagnosis of, 637
epidemiology of, 671
imaging in, 656–658, **729–740**
in elderly persons
assessment of, **573–582**
diagnosis of, 379–380
nail disorders in, 437
onychopathy in, 397
osteoarthritis in, 474–475
pain in, 398–399
patient education on, 574
peripheral neuropathy in, 574–575, 578
preventive care in, **573–582**
surgical management in, 611–612
surgical risks in, 615–616
ulcers in
assessment of, 573–582
in long-term care, 383–384, 387–388, 393–394
pain in, 399
prevention of, 573–582
incidence of, 635
infections in, **655–669**
morbidity and mortality in, 671, 709–710
pathogenesis of, 637–638
peripheral neuropathy in, **671–688**
preoperative considerations with, 215, 219–224
screening for, 637
signs and symptoms of, 638–639
soft tissue reconstruction in, **757–781**
surgery in, **783–792**
treatment of
alpha glucosidase inhibitors in, 643
biguanides in, 641–642
combination, 644
glucose monitoring in, 647
hypoglycemia in, 649
in neuropathy, 677–678
insulin for, 644–647
meglitinides in, 643–644
oral hypoglycemic drugs in, 639–640
perioperative, 647–649
sulfonylureas in, 640–641
thiazolidinediones in, 642
ulcers in, **709–728**
vascular disorders in, **689–708**
versus normal insulin metabolism and glucose homeostasis, 637

Diabetic lumbosacral radiculoplexus neuropathy, 673–674

Dicloxacillin, for diabetic foot infections, 659

Digit(s). *See* Toe(s).

Digital artery island flaps, in diabetic foot reconstruction, 772, 774

Diphenhydramine, for diabetic neuropathy, 679

Disease-modifying antirheumatic drugs, for rheumatoid arthritis, 479–480

Distraction
- arthrodiastasis, 90–92
- in forefoot applications, external fixation for, 36–40

Diuretics, preoperative evaluation of, 226

Doppler studies, in peripheral vascular disease
- in diabetes mellitus, 697–699, 713
- in elderly persons, 539–540

Dornier Epos Ultra device, for plantar fasciitis, 326, 329–330

Dorsalis pedis pulse, evaluation of, in peripheral vascular disease, 533–534

Double-Z rhomboid flaps, in diabetic foot reconstruction, 769–770, 772–773

Dressings
- for diabetic ulcers, 717–718
- for pressure ulcers, 385–387

Drug(s)
- interactions of, in elderly persons, 376
- preoperative evaluation of. *See* Preoperative evaluation, of medications.

Dynafix external fixation system, for forefoot, 27–29

Dynamization, in external fixation, 21

E

Eccrine poroma, 446

Edema, in external fixation, 22

Elderly persons
- arthritis in. *See* Arthritis, in elderly persons.
- atypical illness presentation in, 375–376
- care of, principles of, **373–381**
- deformities in, **487–509, 511–526**
- demographics of, 373
- diabetes mellitus in. *See* Diabetes mellitus, in elderly persons.
- external fixation in, 22
- footwear for, 387, 576, **593–605**
- heterogeneity of, 374
- hyperkeratotic lesions in. *See* Hyperkeratosis, in elderly persons.
- interrelated health events in, 374
- long-term care of, **383–394,** 585
- multidisciplinary care of, 377–378
- multiple conditions in, 375
- nail disorders in. *See* Nail disorders, in elderly persons.
- neurologic evaluation in, **547–571,** 616
- pain in. *See* Pain, in elderly persons.
- peripheral neuropathy in, 379, 574–575, 578
- peripheral vascular disease in. *See* Peripheral vascular disease, in elderly persons.
- podiatric assessment in, 378–379, 383–389, **407–429**
- polypharmacy in, 376
- preoperative evaluation of, 217
- primary care of, **583–591**
- skin disorders in. *See* Skin disorders, in elderly persons.
- surgical considerations in. *See* Surgery, in elderly persons.
- treatment goals for, 377

Electrohydraulic principle, in shock wave generation, 326

Electromagnetic principle, in shock wave generation, 326

Electrophysiologic studies, in diabetes mellitus, 675–676

Elevation dependency test, in peripheral vascular disease, 534–535

Embolism, pulmonary, in deep venous thrombosis, 270–272, 274

Emotional disorders, manifested in foot, 400

Endocrine disorders, preoperative evaluation of, 215

Endothelial injury, deep venous thrombosis in, 270–271

Endovascular therapy, for vascular disease, in diabetes mellitus, 704

Energy expenditures, after amputation, in diabetes mellitus, 799

Enoxaparin, for deep venous thrombosis, 278

Ephedra, preoperative evaluation of, 234

Epidural analgesia, cryotherapy effects on, 263–264

Epinephrine, in local anesthetics, duration of action with, 245

Epos Ultra device, for plantar fasciitis, 326, 329–330

Equinus deformity
external fixation for, 49
in Charcot foot, 99

Ertapenem, for diabetic foot infections, 662

Exercise, for osteoarthritis, 476

Exercise testing, in peripheral vascular disease, 696

Exostectomy, retrocalcaneal, **361–366**

Exostosis
retrocalcaneal, retrocalcaneal exostectomy for, **361–366**
subungual, 434

Extensor digitorum brevis flap, in diabetic foot reconstruction, 777

External fixation
advantages of, 12–13, 66
complications of, 20–22
in forefoot applications, 40
in rearfoot applications, 92–93
disadvantages of, 13–14
in forefoot applications, 30
in rearfoot applications, 66–67
for Charcot foot reconstruction. *See* Charcot foot and ankle, external fixation for.
for deformity correction, 16–18, 47–49, **119–157**
for subtalar arthrodesis, **181–194**
for trauma, 15–16, 35, 47–49, **159–180**
forefoot applications of, **27–44**
fracture after, 21
history of, **1–8,** 9–11, 119, 125, 127
indications for, 14–20, 46–47, 180
midfoot applications of, **45–63**
monolateral devices for, 11
patient selection for, 22–24
psychologic issues in, 23–24
rearfoot applications of, **97–117**
removal of, 21
ring devices for, 11
self-care in, 22–23
training for, 11–12, 14
types of, 11–12, 127, 129

Extracorporeal shock wave therapy, **323–334**
approved devices for, 326
for nonunion, 332
for plantar fasciitis, 326–332
historical review of, 323–324
physics of, 324
physiologic effects of, 324–325
wave generation in, 325–326

F

Factor V Leiden gene mutation, deep venous thrombosis in, 271

Falls, risk assessment for, in long-term care, 392

Fasciotomy, plantar fasciitis, 328

Femoral pulse, evaluation of, in peripheral vascular disease, 533–534

Femoropopliteal reconstruction and bypass, in diabetes mellitus, 702–703

Fibula, fractures of, joint replacement for, case review of, **335–359**

First ray, ulceration at, in diabetes mellitus, surgery for, 786–787

Flaps, in diabetic foot reconstruction, 21, 761–779
local random, 761–773
advancement, 763
advantages of, 761
anatomic considerations in, 762–763
bilobed, 767–769
definition of, 761
double-Z rhomboid, 769–770, 772–773
rhomboid, 769
rotation, 764–765
transposition, 764–766
types of, 761–762
muscle, 21–23, 775
pedicle, 771, 773–779
digital artery island, 774
instep (medial plantar artery), 774
reverse flow sural artery neurofasciocutaneous, 774–776

Flatfoot, 501–503

Flexor digitorum brevis flap, in diabetic foot reconstruction, 778

Fluoxetine, for diabetic neuropathy, 680–681

Food and Drug Administration, shock wave devices approved by, 326

Footwear
for Charcot foot, 747, 751
for diabetic foot, 722
inspection of, 713
ulcerated, 718
for elderly persons, **593–605**
appropriate for activity, 594
construction of, 594–595
custom-molded, 593, 595, 598, 600
hallux valgus due to, 493–494
in claw toe deformity, 490
in deformities, 523–526
in diabetic ulcers, 387, 576
in hallux rigidus, 492–493
in hallux valgus, 495–497
in hammertoe deformity, 490
in metatarsalgia, 499–500

in pes cavus, 505
in pes planus, 503
in plantar heel pain, 506–507
Medicare coverage of, 593
modifications for, 595–598
selection of, 599–600

Forefoot, external fixation applications in, **27–44**
bone grafts, 36
brachymetatarsal correction, 36–38
complications in, 40
compression, 30–36
devices for, 27–30
distraction, 36–40
first metatarsocuneiform joint arthrodesis, 33–35
first metatarsophalangeal joint arthrodesis, 30–32
fractures, 35
Freiberg's disease, 39–40
hallux limitus/rigidus, 39
management in, 40–41
osteotomy, 35–36
shortened first metatarsal, 38–39
soft tissue contraction, 40

Foreign bodies, in nail plate, 435

Fracture(s)
ankle joint replacement for, case review of, **335–359**
external fixation for
calcaneal, 169–170
Lisfranc's, 47–49
metatarsal, 35, 169–172, 174–175
open, 15–16
phalanges, 170–172, 174–175
pilon, 15
talar, 160, 162–166, 169
tibial, 160, 162–166
nonunion of, extracorporeal shock wave therapy for, 332

Freiberg's disease, external fixation for, 39–40

Fungal infections, in elderly persons, 466

G

Gabapentin, for diabetic neuropathy, 681–682

Gait, assessment of
in biomechanical evaluation, 517–518
in neurological evaluation, 551–552

Garlic, preoperative evaluation of, 233

Geriatric patients. *See* Elderly persons.

Ginkgo, preoperative evaluation of, 233

Ginseng, preoperative evaluation of, 233

Glimepiride, for diabetes mellitus, 220, 641

Glipizide, for diabetes mellitus, 220, 641, 644

Glitazones, for diabetes mellitus, 642

Glucosamine, for osteoarthritis, 476

Glucose
homeostasis of, 637
monitoring of, in diabetes mellitus, 647

Glucosidase inhibitors, for diabetes mellitus, 221–222, 643

Glyburide, for diabetes mellitus, 220, 641, 644

Gosnell Scale, for pressure ulcer staging, 385

Gout, in elderly persons, 398, 481–483

Graphesthesia, assessment of, 558

Growth factors, for diabetic ulcers, 717

Guanabenz, preoperative evaluation of, 225

Guanfacine, preoperative evaluation of, 225

Gustillo and Anderson classification, of foot and ankle trauma, 298

H

Haglund's deformity, retrocalcaneal exostectomy for, **361–366**

Half-pin fixator, for tibiotalar arthrodesis, 82, 84–85

Hallux, arthroplasty of, in diabetes mellitus, 786–787

Hallux limitus, 39, 398

Hallux rigidus, 398, 491–493
external fixation for, 39

Hallux valgus, 493–497
biomechanics of, 472
causes of, 493–495
pain in, 398
surgery for, 609
treatment of, 495–497

Hammertoes, 487–491, 785–786

Hapalonychia, 436

Hard corn (heloma durum), 445

Hawkins classification, of foot and ankle trauma, 298

Healing
in elderly persons
factors affecting, 457–458
nutrition for, 458–459, 532, 620–621
of surgical wounds, 616, 622–623
of diabetic ulcers, 715–719, 758

HealthTronics Ossatron device, for plantar fasciitis, 326

Heel
in footwear, modifications of, 597
reconstruction of, in diabetes mellitus, 789–790

Heel pain
plantar, 505–508. *See also* Plantar fasciitis.
retrocalcaneal exostectomy for, **361–366**

Heloma
pain in, 397–398
subungual, 434, 445

Heloma durum (hard corn), 445

Heloma miliare (seed corn), 445

Heloma molle (soft corn), 446

Heloma neurofibrosum, 446

Heloma vasculare, 446

Heparin
for deep venous thrombosis, 277–278
preoperative evaluation of, 228–229

Herbal medications, preoperative evaluation of, 232–234

Hindfoot. *See* Rearfoot.

Hirudin, for deep venous thrombosis, 279

Hives, 460

Hoffman, Raoul, external fixation devices of, 4, 11

Home care, foot care in, 393–394

Hyaluronic acid products, for osteoarthritis, 477

Hydrocodone, for diabetic neuropathy, 680

Hydromorphone, for trauma, 299

Hyperbaric oxygen therapy, for diabetic ulcers, 719

Hypercoagulable states, deep venous thrombosis in, 271, 280

Hyperglycemia, in diabetes mellitus, 639
polyneuropathy development in, 674, 676–677

Hyperkeratosis, in elderly persons, **443–452**
biomechanics of, 444–445
causes of, 444–445
classification of, 443–444
diagnosis of, 447
pain in, 397–398
patient education on, 448
subungual, 434, 437
symptoms of, 445–447
treatment of, 447–448, 584

Hypermobility, of joints, 473–474

Hyperpigmentation, of nail plate, 437

Hypertension, surgical risks in, 615

Hyperuricemia, gout in, 481–483

Hypoglycemia, in diabetes mellitus, treatment of, 649

I

Ibuprofen, for diabetic neuropathy, 679

Ilizarov, Gavriel A., external fixation devices of, 6–8, 77–82, 119, 125, 127

Imaging. *See specific modalities.*

Imipenem, for diabetic foot infections, 664

Imipramine, for diabetic neuropathy, 680–681

Immobilization, in Charcot foot, 747

Immunosuppressive therapy, for rheumatoid arthritis, 480

Infections
in ankle trauma, 294
in diabetes mellitus, **655–669**
antibiotics for, 658–665
costs of, 666
culture in, 656, 719
evaluation of, 655–656
imaging in, 656–658, 729–740
in elderly persons, 577–578
in ulcers, 714
laboratory testing in, 656
pathogens causing, 658
recommendations for, 665–666
surgery for, 665
treatment of, 718–719, 795–796
in elderly persons, 466
in diabetes mellitus, 577–578
of nails, 432
of pressure ulcers, 387
in external fixation, 13, 21, 93, 176–177
local anesthetic failure in, 245

Inflammation
cryotherapy effects on, 309
of nails, 432

Informed consent, for surgery, in elderly persons, 621

Ingrown nails, in diabetes mellitus, 785–786

Injury. *See* Trauma.

Instep (medial plantar artery) flaps, in diabetic foot reconstruction, 774

Insulin
for diabetes mellitus
perioperative, 648–649
regimens for, 646–647
types of, 644–646
metabolism of, 637
preoperative evaluation of, 222–224

Insulin secretagogues, preoperative evaluation of, 221

Intermittent claudication, in peripheral vascular disease, 527–529

Internal fixation
of ankle fractures, arthrosis after, joint replacement for, **335–359**
versus external fixation, 13–14

Interphalangeal joint(s)
first, arthroplasty of, in diabetes mellitus, 786
hammertoe deformity involving, 487–491

Intramedullary nails, for Charcot foot, 751

Irrigation, wound, in trauma, 299–300

Ischemic vascular disease. *See* Peripheral vascular disease.

Island flaps, in diabetic foot reconstruction, 772, 774

J

Joint(s)
Charcot. *See* Charcot foot and ankle.
subluxation of, in external fixation, 22

K

Kaposi's sarcoma, 463

Kava, preoperative evaluation of, 234

Keller arthroplasty, in diabetes mellitus, 787

Ketoacidosis, in diabetes mellitus, 639

Ketorolac, for perioperative pain, 260

Kidney function, preoperative evaluation of, 215

Koilonychia, 437

L

Lambotte, Albin, external fixation devices of, 2–3, 10

Lamotrigine, for diabetic neuropathy, 682

Lasegue's sign, 562

Lauge-Hansen classification, of foot and ankle trauma, 298

Leg, lower, deformities of. *See* Lower leg deformities.

Lepirudin, for deep venous thrombosis, 278–279

Leukocyte scan, in diabetic foot, 735–737

Leukonychia, 436

Levofloxacin, for diabetic foot infections, 659–660

Lidocaine
for diabetic neuropathy, 682
for preemptive analgesia, 246

Limb-length discrepancy
external fixation for, 20
footwear for, 597–598

Limb pressure measurements, in vascular disease, 693–695

Limb salvage
in diabetic foot, 794–797
in severe trauma, versus amputation, 301

Limberg (rhomboid) flaps, in diabetic foot reconstruction, 769

Linezolid, for diabetic foot infections, 659, 661

Lipid-lowering agents, preoperative evaluation of, 226

Lisfranc's fracture, malunion in, external fixation for, 47–49

Lithotripsy. *See* Extracorporeal shock wave therapy.

Liver function, preoperative evaluation of, 215

Local anesthesia, for preemptive analgesia, 243–248

Local random flaps, in diabetic foot reconstruction, 761–773

Long-term care, **383–394**
accidents in, 392
comprehensive plans for, 389
diabetic ulcers in, 387–388, 393–394
in home, 393–394
in nursing homes, 388–389
nutrition in, 391–392
pressure ulcers in, 384–387, 389–391
primary podiatric care in, 585–587
types of, 383

Lower leg deformities, external fixation correction of
anatomic considerations in, 120–124, 126, 129–136

devices for, 127, 129
malalignment evaluation in, 141–143
planning for, 136–143
rotational, 146–147, 150–154
short tibia, 144–145, 148–150
tibial angulation, 133, 145, 148
versus normal values, 137–141

Lower motor neuron lesions, assessment of, 562–563

LY333531, for diabetic neuropathy, 677

M

McGill Pain Questionnaire, 258

Macronychia, 432

Magnetic resonance imaging, in diabetic foot
in vascular evaluation, 701–702
with osteoarthropathy, 657, 737–738

Mal perforans ulcer. *See* Ulcer(s), in diabetes mellitus.

Malgaigne, Jean-Francois, external fixation devices of, *1*, *10*

Malnutrition
perioperative considerations in, 621
skin disorders in, 458–459

Malunion, external fixation for, 18–19, 47–49

Mangled Extremity Severity Score, 301

Medial plantar artery (instep) flaps, in diabetic foot reconstruction, 774

Medicamentosa dermatitis, 460

Medicare
class findings in, 414–415
codes of, 408–411
footwear provisions of, 593

Meglitinides, for diabetes mellitus, 643–644

Melatonin, for diabetic neuropathy, 679

Meperidine, for perioperative pain, 261

Meropenem, for diabetic foot infections, 664

Metabolic disorders, in diabetes mellitus, polyneuropathy in, 676–677

Metastasis, from cancer, 464

Metatarsal(s)
fractures of, external fixation for, 35, 168–170, 172–173
lesser, ulceration at, in diabetes mellitus, surgery for, 788–789
short, correction of, external fixation for, 36–39

Metatarsal bar, in footwear, 596–597

Metatarsalgia, 498–501

Metatarsocuneiform joint, first, arthrodesis of, external fixation for, 33–35

Metatarsophalangeal joint(s)
claw toe deformity involving, 489–491
first
arthrodesis of, external fixation for, 30–32
arthroplasty of, in diabetes mellitus, 786–787
hammertoe deformity involving, 487–491
lateral angulation of. *See* Hallux valgus.
rigidity of, 398, 491–493

Metformin, for diabetes mellitus, 222, 641–642, 644

N-Methyl-D-aspartate receptor antagonists, for preemptive analgesia, 250–251

Metronidazole, for diabetic foot infections, 662–664

Mexiletine, for diabetic neuropathy, 682

Micronychia, 432

Midfoot
external fixation applications in, **45–63**
angular deformity, 47–49
calcaneocuboid arthrodesis, 54–56
cavus osteotomy, *58–59*
Charcot reconstruction. *See* Charcot foot and ankle.
devices for, 46
indications for, 46–47
naviculocuneiform arthrodesis, 57–58
osteomyelitis, 49–52, 59
talonavicular arthrodesis, 56–57
tarsometatarsal joint realignment and arthrodesis, 52–54
reconstruction of, in diabetes mellitus, 789

Miglitol, for diabetes mellitus, 221–222, 643

Minimum Data Set, for nursing home resident assessment, 388–390

Morphine
for perioperative pain, preemptive administration of, 249
for trauma, 299

Morton's neuroma, assessment of, 562

Motor function, assessment of, in elderly persons, 562–566

Motor neuropathy, in diabetes mellitus, 673

Multidisciplinary care, for elderly persons, 377–378

Muscle(s), evaluation of, in motor function assessment, 563–565

Muscle flaps, in diabetic foot reconstruction, 21–23, 775

N

Nafcillin, for diabetic foot infections, 659

Nail disorders
in diabetes mellitus
in elderly persons, 437
surgical treatment of, 785–786
in elderly persons, **431–442**
anonychia, 431
atrophy, 435–436
avulsion, 434–435
Beau's lines, 436
breaking, 437
congenital, 432
contusions, 434
dryness, 437
dystrophy, 436
exostosis, 434
foreign bodies, 435
growth changes, 433
heloma, 434, 445
hemorrhage, 434–435
hyperkeratosis, 434, 437
hyperpigmentation, 437
in diabetes mellitus, 437
in systemic diseases, 431–432, 437–438
inflammatory, 432
ingrown, 435
onycholysis, 431–432, 434–437
onychomycosis, 397, 433–434
onychophosis, 446
pain in, 397
psoriasis, 437
pterygium, 436–437
separation, 436
softening, 436
thickening, 435
treatment of, 585
warts, 432–433

Nateglinide, for diabetes mellitus, 221, 643–644

Naviculocuneiform joint, arthrodesis of, external fixation for, 57–58

Negative pressure therapy, for diabetic ulcers, 718

Neoplasia, of skin, in elderly persons, 463–464

Nerve(s)
cryotherapy effects on, 309
entrapment of, assessment of, 561–562
injury of, in external fixation, 92–93

Nerve blocks, for perioperative pain, 262–263
preemptive, 243–248
agents for, 246–247
duration of action of, 245
factors affecting, 244–245
mechanism of action of, 243–244
peripheral, 247

Nerve conduction studies, in diabetes mellitus, 675

Neuroarthropathy, Charcot. *See* Charcot foot and ankle.

Neurological evaluation
in ankle trauma, 296–297
in diabetes mellitus, 675, 713
of elderly persons, **547–571**, 616
formal examination in, 553–555
history in, 548–550
integrated sensorimotor function, 566–569
motor function, 562–566
observation in, 550–553
sensory function, 555–562
terminology of, 569–570

Neuroma, Morton's, assessment of, 562

Neuropathy. *See also* Peripheral neuropathy.
Charcot. *See* Charcot foot and ankle.
pain in, 399–400

Nikolsky's sign, 465

NMDA (N-methyl-D-aspartate) receptor antagonists, for preemptive analgesia, 250–251

Noncontact normothermic wound therapy, for diabetic ulcers, 718

Nonsteroidal anti-inflammatory drugs
for gout, 482
for osteoarthritis, 477
for perioperative pain, 250, 259–260
for rheumatoid arthritis, 479
preoperative evaluation of, 232

Nonunion
external fixation for, 18–19
extracorporeal shock wave therapy for, 332
in external fixation, 22

Norton Scale, for pressure ulcer staging, 385

Nortriptyline, for diabetic neuropathy, 680

NSAIDs. *See* Nonsteroidal anti-inflammatory drugs.

Numbness, in diabetic neuropathy, 673–674

Nursing homes, foot care in, 388–393

Nutrition
  assessment of, in amputation planning, 796
  for elderly persons, 458–459, 532
    assessment of, 549
    in long-term care, 391–392
    surgical considerations in, 620–621

O

Obesity
  in osteoarthritis, 476
  preoperative evaluation of, 218

Occlusive vascular disease. *See* Arterial occlusive disease; Peripheral vascular disease.

Occupational disorders
  neurologic, 549
  skin, 460

Ofloxacin, for diabetic foot infections, 659–660, 664

Onychatrophia, 435–436

Onychauxis, 397, 432

Onychexallis, 433

Onychia, 432

Onychia punctata, 437

Onychia sicca, 437

Onychochauxis, 435

Onychoclasis, 437

Onychocryptosis, 435

Onychodystrophy, 436

Onychogryphosis, 397, 432

Onycholysis, 431–432, 434–437

Onychomadesis, 434–436

Onychomalacia, 434, 436

Onychomycosis, 397, 433–434

Onychophosis, 397, 434, 437, 446

Onychophyma, 433

Onychoptosis, 437

Onychorrhexis, 436

Onychoschizia, 436

Onychyphemia, 434–435

Open fractures, external fixation for, 15–16

Open reduction, in Charcot foot, 749, 751

Opioids
  for diabetic neuropathy, 679–680
  for perioperative pain, 249, 260–262

Oral antidiabetic agents, 220–222, 639–644

Orthofix external fixation system, for forefoot, 27–29

Orthoses. *See also* Footwear.
  for Charcot foot, 751
  for claw toes, 490
  for deformities, 523–526
  for diabetes mellitus, 576
  for hallux rigidus, 492–493
  for hallux valgus, 495–497
  for hammertoes, 490
  for metatarsalgia, 500–501
  for pes cavus, 505
  for pes planus, 503
  for plantar heel pain, 507–508
  for tailor's bunion, 497–498

Ossatron device, for plantar fasciitis, 326

Osteoarthritis
  in elderly persons
    biomechanics of, 470–472
    epidemiology of, 469–470
    joints affected by, 472
    pain in, 398
    pathogenesis of, 474–475
    surgery for, 609–610
    treatment of, 472–477
  post-traumatic, ankle joint replacement for, case review of, **335–359**
  synovectomy for, external fixation after, 16

Osteoarthropathy, Charcot. *See* Charcot foot and ankle.

Osteomyelitis
  in diabetes mellitus
    antibiotics for, 658–665
    imaging in, 656–658, 729–740
    of lesser metatarsals, 788
    pathogens causing, 658
    recommendations for, 665–666
    surgery for, 665
    treatment of, 718–719
    versus Charcot foot, 746
  in elderly persons
    in pressure ulcers, 386
    in subungual abscess, 432
    pain in, 399
  midfoot, external fixation for, 49–52, 59

Osteophytes, in osteoarthritis, 475

Osteoporosis
  pain in, 398
  preoperative considerations in, 613

Osteotomy
for cavus foot, external fixation for, 58–59
of metatarsals
external fixation of, 35–36
in diabetes mellitus, 788–789

Oxacillin, for diabetic foot infections, 659

Oximetry, in peripheral vascular disease, 539, 696–697

Oxycodone
for diabetic neuropathy, 679–680
for perioperative pain, 261

Oxygen therapy, for diabetic ulcers, 719

P

Pachonychia, 432

Pain. *See also* Pain management.
assessment of, 257–259
heel
in plantar fasciitis. *See* Plantar fasciitis.
retrocalcaneal exostectomy for, **361–366**
in arterial occlusive disease, 690
in diabetic neuropathy, 673–674, 678–683
in elderly persons, **395–406**
causes of, 395–396
impact of, 395–396
in arterial ulcers, 389
in arthritis, 398
in biomechanical disorders, 398
in diabetes mellitus, 398–399
in gout, 481–482
in hallux rigidus, 492
in hyperkeratosis, 397–398
in nail disorders, 397, 432, 434–435
in neurologic disorders, 399–400
in osteoarthritis, 476–477
in peripheral vascular disease, 399, 527–530
in pes cavus, 504–505
in systemic disease, 398
metatarsalgia, 498–501
perception of, assessment of, 558–559
plantar heel, 505–508
psychosocial correlates of, 400–401
skin changes in, 396–397
in external fixation, 22
physiology of, 237–240, 307–310

Pain management
in diabetic neuropathy, 678–683
in extracorporeal shock wave therapy, for plantar fasciitis, 329
in trauma, 298–299
postoperative, **257–267**
after return to home, 264–265
assessment in, 257–259
cryotherapy for, **307–322**
neural block in, 262–263
NSAIDs in, 259–260
opioids in, 260–261
patient-controlled analgesia in, 262
patient education in, 264
psychosocial issues in, 199–211
spinal/epidural analgesia in, 263–264

Parkhill, Clayton, external fixation devices of, 2, 10

Paronychia, 397

Paroxetine, for diabetic neuropathy, 681

Patient-controlled analgesia, for perioperative pain, 262

Pediatric patients, external fixation in, 22

Pedicle flaps, in diabetic foot reconstruction, 771, 773–779

Pemphigus vulgaris, 464–465

Pennig external fixation system, for forefoot, 27–28

Pentoxifylline, for diabetic neuropathy, 683

Percutaneous transluminal angioplasty, for vascular disease, in diabetes mellitus, 704

Peripheral neuropathy
in diabetes mellitus, **671–688**
amputation risk in, 783–784
classification of, 673–674
clinical course of, 674
definition of, 672
diagnosis of, 674–676
differential diagnosis of, 675–676
epidemiology of, 672–673
in elderly persons, 574–575, 578
pathology of, 676–677
treatment of, 677–683
ulcers in, 711
in elderly persons
diagnosis of, 379
flatfoot in, 502
in diabetes mellitus, 574–575, 578
types of, 554

Peripheral vascular disease. *See also* Arterial occlusive disease.
in elderly persons, **527–545**
diagnosis of, 379
healing ability and, 531–532

history of, 527–532
in diabetes mellitus, 575
noninvasive evaluation of, 539–543
pain in, 399
physical examination in, 532–538
risk factors for, 530–531
surgical risks in, 615–616
venous, 543–545

Peroneal bypass grafting, in diabetes mellitus, 703–704

Pes cavus, 503–505

Pes planus, 501–503

Phalanges, fractures of, external fixation for, 168–170, 172–173

Phenytoin, for diabetic neuropathy, 681

Photoplethysmography, in peripheral vascular disease, 539–541

Physical therapy
for ankle joint replacement, 340–341
for osteoarthritis, 476

Piezoelectric methods, for shock wave generation, 325–326

Pilon fractures, external fixation for, 15, 158, 160–164

Pioglitazone, for diabetes mellitus, 221, 642

Piperacillin, for diabetic foot infections, 664

Piperacillin/tazobactam, for diabetic foot infections, 660, 662–664

Plantar artery flaps, in diabetic foot reconstruction, 774

Plantar fasciitis, 505–508
conservative treatment of, 327–328
extracorporeal shock wave therapy for
adverse effects of, 329
approved devices for, 326
cautions with, 329
clinical studies of, 329–331
indications for, 328
procedure for, 329
fasciotomy for, 328
pathophysiology of, 327

Plantar keratosis, 447

Plate fixation, of ankle fractures, arthrosis after, joint replacement for, **335–359**

Platelet-derived growth factor gel, for diabetic ulcers, 717

Plethysmography, in peripheral vascular disease, 539–541

Plexopathy, 554

Podogeriatric care. *See* Elderly persons.

Polymicrobial infections, in diabetic foot, 658

Polyneuropathy, diabetic, 673–677

Polyol pathway, in diabetes mellitus, 677

Polyonychia, 432

Polypharmacy, in elderly persons, 376

Popliteal pulse, evaluation of, in peripheral vascular disease, 533–534

Porokeratosis plantaris discreta, 447

Poroma, eccrine, 446

Porphyria cutanea tarda, 465

Posterior tibial tendon dysfunction, 502

Postocclusive reactive hyperemia test, in peripheral vascular disease, 535–538

Posture, assessment of
in biomechanical evaluation, 515
in neurological evaluation, 550

Prednisone, preoperative evaluation of, 230

Preemptive analgesia, perioperative, **237–255**
benefits of, 242–243
clinical relevance of, 242–243
definition of, 242
goals of, 240–241
historical perspective of, 241
local, 243–248
misconceptions about, 241–242
multimodal, 251
NMDA receptor antagonists for, 250–251
NSAIDs for, 250
opioids for, 249
pain pathophysiology and, 237–240
preincisional versus postincisional block, 248
presurgical versus postsurgical block, 248

Preoperative evaluation, **213–235**
for diabetic foot elective procedures, 784–785
history-taking in, 213–217
laboratory testing in, 216–217
of medications, 219–234
anticoagulant, 226–229
antidepressant, 231
antidiabetic, 219–224
antiplatelet, 230
antipsychotic, 231
anxiolytic, 231
aspirin, 231–232
cardiovascular, 224–226

complementary, 232–234
corticosteroids, 230
NSAIDs, 231–232
risk factor assessment in, 217–219

Pressure relief, for diabetic ulcers, 718

Pressure Specifying Sensory Device, for sensory evaluation, 557–558

Pressure ulcers
in long-term care, 384–387, 389–391
assessment of, 384–385
prevention of, 389–390
risk for, 385, 390
staging of, 384–386
treatment of, 386–387, 391
postoperative, 624

Primary podiatric medicine, **583–591**
assessment in, 584
goals of, 586
in long-term care settings, 585–587
program for, 586–587
settings for, 584
standards of care in, 584
treatment in, 584–585

Probenecid, for diabetic foot infections, 659

Pronation
arthritis development in, 471
orthoses for, 525

Propranolol, preoperative evaluation of, 224–225

Proprioception, evaluation of, 559

Protein C, activated, resistance to, deep venous thrombosis in, 271

Protein S, defects of, deep venous thrombosis in, 271

Prothrombin, defects of, deep venous thrombosis in, 271

Psoriasis, 462–463

Psychiatric disorders, foot pain in, 401

Psychologic issues
in external fixation, 23–24
in foot pain, 400–401

Psychosocial management, of surgery patient, **199–211**
high-risk, 204
pretreatment screening, 200–201, 215
risk factor assessment and application in, 204–208
surgical outcome and, 201–204

Pterygium, of nail wall, 436–437

Pulmonary embolism, in deep venous thrombosis, 270–272, 274

Pulmonary function testing, preoperative, 214

Pulse, evaluation of
in diabetes mellitus, 713
in peripheral vascular disease, 533–534

Pulse volume recording, in peripheral vascular disease, 539, 695–696

Pump bump (Haglund's deformity), retrocalcaneal exostectomy for, **361–366**

Q

Quality of life, in foot pain, 400–401

Quantitative sensory testing, in diabetes mellitus, 675

Quinupristin-dalfopristin, for diabetic foot infections, 659, 662

R

Racial factors, in amputation, in diabetes mellitus, 799–800

Radiculopathy, 555

Radiography
in ankle trauma, 297–298
in diabetic foot, 656–657, 730–732

Ram's horn nails, 432

Random flaps, local, in diabetic foot reconstruction, 761–773

Rearfoot
alignment of, assessment of, 515–516
external fixation applications in, **65–96**
biomechanics of, 67–69
capabilities of, 66
circular, 67–69
complications of, 92–93
distraction arthrodiastasis, 90–92
frame removal in, 94
limitations of, 66–67
tibiocalcaneal arthrodesis, 16, 85–89
tibiotalar arthrodesis, 16, 74–85
triple arthrodesis, 16, 89–90
wire considerations in, 69–74
reconstruction of, in diabetes mellitus, 789–790

Reflexes, assessment of, 553, 566–568

Repaglinide, for diabetes mellitus, 221, 643–644

Resident Assessment Protocols, for nursing home residents, 388

Rest pain, in peripheral vascular disease, 530

Retrocalcaneal exostectomy, **361–366**

Reverse flow sural artery neurofasciocutaneous flaps, in diabetic foot reconstruction, 774–776

Rheumatoid arthritis
in elderly persons, 477–481
biomechanics of, 470–472
diagnosis of, 477–478
epidemiology of, 469
joints affected by, 472
pain in, 398
treatment of, 472–474, 478–481
subtalar arthrodesis in, external fixation for, 183–188

Rhomboid flaps, in diabetic foot reconstruction, 769–770, 772–773

Rifampicin, for diabetic foot infections, 660

Rigidity, in motor function assessment, 564

Risk factors
for deep venous thrombosis, 269–270, 272–273
preoperative evaluation of, 217–219
psychosocial, assessment of, presurgical, 204–208

Road to Recovery pain assessment tool, 258–259

Rocker bar, in footwear, 597

Rockerbottom foot, external fixation for, 49–52

Rofecoxib, for perioperative pain, 250, 260

Romberg's test, 568–569

Ropivacaine, for preemptive analgesia, 246

Rosiglitazone, for diabetes mellitus, 221, 642, 644

Rotation flaps, in diabetic foot reconstruction, 764–765

S

Sagittal deformity, of midfoot, external fixation for, 49

Saint-John's wort, preoperative evaluation of, 233

Saline dressings, for diabetic ulcers, 717

Scintigraphy, in diabetic foot, 732–735
with osteoarthropathy, 657
with ulcers, 714–715

Screw fixation
of ankle fractures, arthrosis after, joint replacement for, **335–359**
of first metatarsocuneiform joint arthrodesis, external fixation with, 33–35

Sedation, in trauma, management of, 298–299

Seed corn (heloma miliare), 445

Segmental limb pressure measurements, in vascular disease, 693–695

Semmes-Weinstein monofilaments, for sensory evaluation, 556–557, 675, 713

Sensorimotor function, integrated, assessment of, 566–569

Sensory function, evaluation of
in diabetes mellitus, 713
in elderly persons, 555–562

Sensory neuropathy, in diabetes mellitus, 673

Sesamoidectomy, in diabetes mellitus, 787

Shank filler, in footwear, 597

Shock wave therapy, extracorporeal. *See* Extracorporeal shock wave therapy.

Shoes. *See* Footwear.

Skewfoot, external fixation for, 48–49

Skin, temperature of, in arterial occlusive disease, 691–692

Skin disorders
in diabetes mellitus, examination of, 713–714
in elderly persons, **453–467**
aging changes and, 454, 459
biopsy in, 454–456
blistering, 464–465
contact dermatitis, 460
diagnosis of, 454–455
in diabetes mellitus, 574–575
in malnutrition, 458–459
in peripheral vascular disease, 532–533
in systemic disease, 453
neoplastic, 463–464
pain in, 396–397
psoriasis, 462–463
terminology of, 454–459
ulcers. *See* Ulcer(s).
warts, 432–433, 446–447, 464
xerosis, 460, 462

Skin grafting, for diabetic foot, 759–761

Socks, for metatarsalgia, 499–500

Soft corn (heloma molle), 446

Soft tissue
contraction of, prevention of, external fixation in, 40
injury of, in external fixation, 21
reconstruction of, in diabetic foot, **757–781**
anatomic considerations in, 757–758
local random flaps in, 761–773
pedicle flaps in, 771, 773–779
skin grafting in, 759–761
team approach to, 757

Solid ankle cushion heel, in footwear, 597

Spasticity, in motor function assessment, 564

Spinal analgesia, cryotherapy effects on, 263–264

Spoon nails, 437

Staphylococcal infections, in diabetic foot, 658, 660–661

Stereognosia, assessment of, 558

Straight leg raising test, 562

Strength testing, in neurological evaluation, 564–566

Streptococcal infections, in diabetic foot, 658, 660–661

Stress, in hyperkeratosis formation, 444–445

Subpapillary venous plexus filling time test, in peripheral vascular disease, 534

Subtalar joint, arthrodesis of, external fixation for, **181–194**
case studies of, 185–194
in joint degeneration, 190–193
in rheumatoid arthritis, 185–190
indications for, 180
postoperative management in, 185
technique for, 182–185

Sudomotor axon reflex test, in diabetes mellitus, 675

Sulfonylureas, for diabetes mellitus, 220, 640–641, 644

Sulindac, for diabetic neuropathy, 679

Sural artery flaps, in diabetic foot reconstruction, 774–776

Surgery. *See also specific disorders and procedures.*
ankle joint replacement, **335–359**
cryotherapy after, **307–322**
deep venous thrombosis in. *See* Deep venous thrombosis.
extracorporeal shock wave, for plantar fasciitis, **323–334**
for diabetic foot
disease management in, 647–649
elective, **783–792**
Achilles tendon lengthening, 790
first ray, 786–787
goals of, 784
lesser digits, 785–786
lesser metatarsals, 788–789
midfoot, 789
nails, 785–786
preoperative evaluation for, 784–785
rearfoot, 789–790
in infections, 662–665
in ulceration, 720–721
for trauma, **291–306**
in elderly persons, **607–626**
age considerations in, 607
anesthesia for, 607
complications of, 623–624
for gout, 483
for osteoarthritis, 477
for rheumatoid arthritis, 480–481
goals of, 612
in peripheral vascular disease, 541–543
in primary care setting, 585
indications for, 608–612
perioperative care in, 621–622
postoperative care in, 622–623
preoperative assessment in, 619–621
risks of, 614–619
systems review for, 613–614
pain control in. *See* Pain management, postoperative.
preemptive analgesia for, **237–255**
preoperative evaluation and management for, **213–235**
psychosocial management in, **199–211**
retrocalcaneal exostectomy, **361–366**

Swelling, cryotherapy effects on, 309

Syme amputation, in diabetic foot, 797–798

Synovectomy, for osteoarthritis, external fixation after, 16

Synthes tubular external fixation, for subtalar arthrodesis, **181–194**
case studies of, 185–193
in joint degeneration, 191–193
in rheumatoid arthritis, 185–190
indications for, 182
postoperative management in, 185
technique for, 182–185

T

Tactile Circumferential Discriminator, for sensory evaluation, 557

Tailor's bunion, in elderly persons, 497–498

Talonavicular joint, arthrodesis of, external fixation for, 56–57

Talus, fractures of, external fixation for, 160, 162–166, 169

Tarsometatarsal joint, arthrodesis of, external fixation for, 52–54

Temperature sensation, assessment of, 559–560

Terry's nails, 437

Tetanus, in ankle trauma, 294–295

Thiazolidinediones, for diabetes mellitus, 221, 642

Thomas heel, in footwear, 597

Thrombosis, venous. *See also* Deep venous thrombosis.
in elderly persons, 544–545, 616

Tibia
angulation of, external fixation correction of, 133, 145, 148
deformities of. *See* Lower leg deformities.
fractures of, joint replacement for, case review of, **335–359**
pilon fractures of, external fixation for, 15, 160, 162–166
rotational deformity of, external fixation correction of, 146–147, 150–154
short, external fixation correction of, 144–145, 148–150

Tibial pulse, posterior, evaluation of, in peripheral vascular disease, 533–534

Tibiocalcaneal joint, arthrodesis of, external fixation for, 16, 85–89

Tibiotalar joint, arthrodesis of, external fixation for, 16, 74–85

Ticarcillin/clavulanate, for diabetic foot infections, 662–664

Ticlopidine, preoperative evaluation of, 230

Tinazaprin, for deep venous thrombosis, 278

Tinea unguium, pain in, 397

Tinel's sign, 562

Toe(s)
alignment of, assessment of, 516–517
claw, 489–491
deformities of, in diabetes mellitus, surgical treatment of, 785–786
gout in, 398, 481–483
great. *See subjects starting with* Hallux.
hammer deformity of, 487–491, 785–786
heloma of. *See* Heloma.
pressure measurements in, in vascular disease, 695
tailor's bunion of, 497–498

Tolazamide, for diabetes mellitus, 640–641

Tolbutamide, for diabetes mellitus, 640–641

Topiramate, for diabetic neuropathy, 682

Total-contact cast, for diabetic ulcers, 718

Touch, evaluation of, 556–558

Tourniquet, for trauma repair, 301

Tramadol, for diabetic neuropathy, 680

Trandolapril, for diabetic neuropathy, 683

Transcutaneous electrical nerve stimulation, for diabetic neuropathy, 683

Transcutaneous oximetry, in peripheral vascular disease, 539, 696–697

Transposition flaps, in diabetic foot reconstruction, 764–766

Trauma. *See also* Fracture(s).
arthrosis after, ankle joint replacement for, case review of, **335–359**
external fixation for, **159–180**
ankle arthrodesis, 166–169
ankle joint bridging, 160, 162–166
calcaneal fractures, 169–170
contractures, 176, 178
Ilizarov technique for, 172–173, 176–179
metatarsal fractures, 170–172, 174–175
phalangeal fractures, 169–172, 174–175
talar fractures, 169
in diabetes mellitus, ulcers in, 711
perioperative management of, **291–306**
dressings in, 303–304
historical perspective of, 292–293
history taking in, 294–295
in displaced fracture, 302
in open injury, 299–302
initial evaluation in, 293
internal fixation in, 302–303
pain control in, 298–299
physical examination in, 295–298
surgical procedure in, 299–305
systematic evaluation in, 294–298
wound closure in, 303

Tremor, assessment of, in neurological evaluation, 550–551

Tricyclic antidepressants
for diabetic neuropathy, 680–681
preoperative evaluation of, 231

Triple arthrodesis, external fixation for, 16, 89–90

Tuning fork, for vibratory sensation assessment, 560–561

Two-point discrimination test, for sensory evaluation, 557–558

Tyloma (callus), 397–398, 445

U

Ulcer(s)
in diabetes mellitus, **709–728**. *See also* Ulcer(s), in elderly persons, in diabetes mellitus.
classification of, 715
epidemiology of, 710
etiology of, 711–712
evaluation of, 712–715
prevention of, 721–722. *See also* Surgery, for diabetic foot, elective.
risk factors for, 783–784
treatment of
healing promotion modalities in, 715–719
surgical, 720–721
in elderly persons, 465–466
arterial, 389
in diabetes mellitus
assessment of, 573–582
in long-term care, 387–388, 393–394
pain in, 399
prevention of, 573–582
pressure, in long-term care, 384–387, 389–391
treatment of, 466
venous stasis, 388–389
pressure. *See* Pressure ulcers.

Ultrasonography
Duplex
in deep venous thrombosis, 274–275
in peripheral vascular disease, 697–699
in Charcot foot, in diabetes mellitus, 657–658

University of Texas, wound classification system of, 715

Upper motor neuron lesions, assessment of, 563

Urate deposition, in gout, 481–483

Urticaria, 460

V

V-Y advancement flap, in diabetic foot reconstruction, 763–764

Vacuum-assisted closure, of diabetic ulcers, 718

Valerian, preoperative evaluation of, 234

Valgus deformity, of midfoot, external fixation for, 48–49

Valproate, for diabetic neuropathy, 683

Vancomycin, for diabetic foot infections, 659–661, 664

Varus deformity, of midfoot, external fixation for, 48–49

Vascular examination, in ankle trauma, 295–296

Vasculopathy, in diabetes mellitus
polyneuropathy in, 676–677
ulcers in, 711

Vena cava, inferior, interruption therapy for, for deep venous thrombosis, 280

Venography, in deep venous thrombosis, 275

Venous disease, in elderly persons, 543–545
stasis ulcers, 388–389
surgical risks in, 616

Venous stasis, deep venous thrombosis in, 270–271

Venous thrombosis, deep. *See* Deep venous thrombosis.

Verrucae. *See* Warts.

Vibratory sensation, assessment of, 560–561

W

Wagner Scale, for diabetic ulcer staging, 387

Warfarin
for deep venous thrombosis, 279
preoperative evaluation of, 227–228

Warts, in elderly persons, 446–447
of nail area, 432–433
treatment of, 464

Wedges, in footwear, 595–596

Whartenberg pinwheel, in pain perception assessment, 558

Wires, in external fixation
of first metatarsocuneiform joint arthrodesis, 33–35
of first metatarsophalangeal joint arthrodesis, 30–32
of rearfoot procedures, 69–74

Wound. *See also* Ulcer(s).
assessment of, in diabetic ulcers, 714–715
treatment of, in pressure ulcers, 386–387

X

Xerosis, 460, 462

Z

Z-plasty, in double-Z rhomboid flaps, in diabetic foot reconstruction, 769–770, 772–773

Zolpidem, for diabetic neuropathy, 679